

## Does Improvement in Patient Pain and Fatigue Lag Behind Clinical Remission in Rheumatoid Arthritis Patients? Data from a Rheumatoid Arthritis Registry

Janet Pope, Emmanouil Rampakakis, Mohammad Movahedi, Angela Cesta, John Sampalis and Claire Bombardier

**Background/Purpose:** Rheumatoid arthritis (RA) patients are often not in remission due to patient global assessment of disease activity (PtGA) which is included in the formula of all disease activity indices. Given that PtGA may reflect pain or fatigue, the aim of this analysis was to assess the relative timing and perhaps lag of patient-reported outcomes (PRO) after remission is obtained as measured by clinical disease activity index (CDAI) or swollen joint count (SJC28) in a large observational database of RA patients followed in routine clinical care.

**Methods:** RA patients enrolled in the Ontario Best Practices Research Initiative (OBRI) registry that were not in low disease state at baseline based on the CDAI, SJC28, PtGA, pain and fatigue criteria below, and had at least six months of follow-up, were included in the analysis. Low disease state was defined as  $CDAI \leq 10$ ,  $SJC28 \leq 2$ ,  $PtGA \leq 2$ cm, pain score  $\leq 2$ cm, or fatigue score  $\leq 2$ cm. Remission was defined as  $CDAI \leq 2.8$ ,  $SJC28 \leq 1$ ,  $PtGA \leq 1$ cm, pain score  $\leq 1$ cm, or fatigue score  $\leq 1$ cm. Kaplan-Meier survival analysis was used to assess the time to first low disease state / remission based on each definition.

**Results:** A total of 986 patients were included with a mean (SD) age and disease duration of 57.4 (12.9) years and 8.3 (9.9) years, respectively, and mostly women (80.0%). Mean (SD) of CDAI, SJC28, PtGA, pain, and fatigue at enrolment was 29.8 (11.7), 8.3 (4.6), 6.4 (1.9), 6.6 (1.9), and 6.7 (2.0), respectively.



Figure 1 shows the time to first low disease state and time to first remission based on different definitions. The median (95%CI) time in months to  $CDAI \leq 10$  was 12.4 (11.4-13.6),  $SJC28 \leq 2$  was 9 (8.2-10),  $PtGA \leq 2$ cm was 18.9 (16.1-22), pain  $\leq 2$ cm was 24.5 (19.4-30.5), and fatigue  $\leq 2$ cm was 30.4 (24.8-31.7). For remission, the median (95%CI) time to  $CDAI \leq 2.8$  was 46.5 (42-54.1),  $SJC28 \leq 1$  was 12.5 (11.4-13.4),  $PtGA \leq 1$ cm was 39.6 (34.6-44.8), pain  $\leq 1$ cm was 54.7 (43.6-57.5), and fatigue  $\leq 1$ cm was 42.6 (36.8-48).

**Conclusion:** Time to achieving low disease state or remission based on PROs is considerably longer compared to swollen joint count which may have significant impact on the time to CDAI low disease activity and remission. Remission and low disease activity composite scores, and PROs lag behind SJCs. Careful interpretation of PROs and composite scores could impact management including prevention of overtreatment and unnecessary switching of DMARDs.



### A. Time to First LDA

### B. Time to First Remission



CDAI





SJC-28



PtGA



Pain



Fatigue

